



# MESA Exam 6 Plus

Participant ID #:

Acrostic:

Phlebotomist ID:

Date:

/

/

Month

Day

Year

QC ID:


## Urine / Phlebotomy

### PARTICIPANT QUESTIONS


	Yes	No	Don't know
1. ¿Sangra o le aparecen morados o moretones con facilidad?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2. ¿Le han dicho alguna vez que tiene un trastorno de la coagulación de la sangre?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3. ¿Alguna vez se desmayó mientras le extraían sangre?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4. ¿Padece de diabetes tratada con insulina o hipoglucemiantes orales?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

### PROCEDURE

5. Was urine sample filled?

☐ Yes

Skip to #6

☐ No☐ Partial

mL

Why was urine sample not taken?

☐ Participant unable to void☐ Refused☐ Other:

6. Time at start of venipuncture:

:

Hr

Min

☐ AM☐ PM

7. Was any blood drawn?

☐ Yes, full sample☐ Yes, partial sample☐ No, refused☐ No, hard to stick☐ No, other:8. Elapsed time until  
tourniquet released:

seconds

(120-seconds optimum)

9. Time at end of  
venipuncture:

:

Hr

Min

☐ AM☐ PM



## Exam 6 Plus

### Urine / Phlebotomy

10. Quality of venipuncture: ☐ Traumatic ☐ Clean



Mark all  
that apply

☐ Vein collapsed

☐ Excessive duration of draw

☐ Vein hard to get at

☐ Hematoma

☐ Multiple sticks

☐ Leakage at venipuncture site

If tube is not full, but is at least half full, please indicate "Partial" and enter the volume to the nearest mL.

11. Blood volume per tube:

**Filled**

Specify  
volume (mL):  
*min 1/2 full*

Exam 6A:

Yes

No

Partial

a. Serum 7.5 mL

☐

☐

☐

*Include if consented to vitD study*

b. Paxgene 2.5 mL

☐

☐

☐

*Include if consented to vitD study*

Comments: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_